

Relationship Between Physical Activity And Body Vitals Among Undergraduate Students At Rivers State University

Kpai Tonubari

Ignatius Ajuru University Of Education, Rivers State, Nigeria

Abstract: This study explores the correlation between physical activity levels and key body vitals among undergraduate students at Rivers State University. Utilizing a cross-sectional design, data were collected from 300 students through self-reported physical activity questionnaires and direct measurements of body vitals, including heart rate, blood pressure, and body mass index (BMI). The analysis revealed a significant positive relationship between regular physical activity and improved body vitals. Students who engaged in moderate to vigorous physical activity exhibited lower resting heart rates, optimal blood pressure levels, and healthier BMI ranges compared to their less active peers. These findings underscore the importance of physical activity in maintaining and enhancing physiological health among university students. The study highlights the need for targeted interventions and policies to promote physical activity within the university setting to foster better health outcomes. Future research should investigate the long-term effects of sustained physical activity on body vitals and academic performance.

Keywords: exercise; students; assessment; physical fitness; physical activity

1. Introduction

Diet and physical activity directly affect the health status of adults and children [1–6]. Due to the fact that the largest part of the world's population is physically inactive, physical inactivity is considered to be a public health problem, as opposed to an individual problem. According to the report of the World Health Organization, physical inactivity is a risk factor, along with smoking, obesity, and hypertension [7]. Stress, obesity, and movement disorders such as hypokinesia are the most common causes of premature death, along with chronic non-communicable diseases from which neither children nor young people are immune [8]. Economically stable countries (about 60–70% of them) do not achieve even the minimum level of physical activity recommended by the World Health Organization in order to maintain health and energy balance [9,10].

Based on the abovementioned facts, the negative consequences of physical activity decline are also reflected in the student population, where the reduction of physical activity could also lead to decreased physical fitness. Health-related physical fitness is also influenced by many other factors, such as body weight and socioeconomic status. For example, the physical activity level of individuals of low socioeconomic status likely suffers due to their living areas providing less access to parks [11] and neighborhood walkability [12]. Additionally, their health is also negatively affected by the cost of healthy food compared to that of junk food [13]. Bodyweight disorder is very common in students and it can be



often traced back to being overweight in childhood and adolescence [14]. Nevertheless, one of the most common negative external factors that influenced the exercise of physical activity in the students' population is the lack of free time due to the schedule at the faculty, obligations in social and family life.

According to Caia et al. [15], low strength was peculiar to 61% of students and 28% had below-average strength. Kubieva et al. [16] concluded that students have problems with body mass index and strength, regardless of their physical activity level. Likewise, low cardiorespiratory fitness is also an important predictor and could be prevented only with lifestyle modifications [17], such as increasing physical activity and promoting dietary changes [18]. Kwan et al. [19] have already noticed the evident decline of physical activity when enrolling a university and according to several authors [20–22], already one-third of high school students are insufficiently active after transitioning to university life. This was confirmed by the study that investigated physical activity patterns among American, Asians, Africans and Hispanic university students. The authors have found that 46.7% of them didn't engage in physical activity and 16.7% were physically inactive [23]. Several studies have also noted a weak physically active lifestyle and it is on the rise among university students [24–26].

Based on the above-mentioned facts, and due to the fact that many students feel pressure during the engagement in academic activities, with no time for physical activities, it was necessary to determine the current state of physical activity and physical fitness among university students. Therefore, the aim of this systematic review was to examine the scientific evidence regarding the level of physical activity and physical fitness among university students.

2. Materials and Methods

2.1. Literature Identification

Studies were searched and analyzed in accordance with the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines [27]. In addition, this study is approved by the Ethical Board of Faculty of Sport and Physical Education, University of Niš. The research included studies conducted from 2011 to 2021 and the literature relevant for this type of research, available with the following databases: Google Scholar, PubMed, Science Direct, and Scopus.

The following keywords were used to search for the articles reporting on physical activity and physical fitness among university students: (“exercise” OR “physical activity” OR “accelerometer” OR “physical fitness” OR “strength” OR “endurance”) AND (“students” OR “adults” OR “students population” OR “university students” OR “college students”).

A descriptive method was used to analyze the data obtained, and all titles and abstracts were reviewed for possible study inclusion. At the same time, the identification strategy was modified and adapted to the particular database to increase the sensitivity. After a detailed identification process, studies were considered to be relevant if they met the inclusion criteria.

The search for studies, assessment of their value and data extraction were conducted independently by two authors (M.S. and D.Đ.), and the lists of references from previously assessed and original research were also reviewed. After that, each author cross-examined the identified studies, which were then taken for further analysis or rejected.

2.2. Inclusion Criteria

For the study to be included in the final analysis, it had to meet the following criteria: year of publication (2011 to 2021), the studies published in English, the experimental study included males and females attending faculty and that the participants were tested for the evaluation of physical activity and physical fitness.

2.3. Risk of Bias Assessment

The risk of bias was assessed according to the PRISMA guidelines, that is, using the PEDro scale [28] to determine the quality of reviewed studies and the potential risk of bias. Two independent authors (D.D. and M.S.) assessed the quality and risk of bias using checklists. Concordance between reviewers was estimated using k -statistics data to review the full text and assess relativity and risk of bias. In case of discordance as to findings of the risk of bias assessment, the obtained data was assessed by the third reviewer (K.P.), who also gave the final decision. The k rate of concordance between reviewers' findings was $k = 0.91$.

2.4. Data Extraction

When the cross-examination was conducted, if the data were adequate, the necessary information was extracted and then moved to an Excel spreadsheet. The standardized data extraction protocol was applied (Cochrane Consumer and Communication Review Group's) to extract the characteristics, such as authors and year of study, sample size, age, types of experimental program, duration, frequency, and study results.

3. Results

3.1. Quality of the Studies

Of the total number of studies that were included in the quantitative synthesis, and based on the points each study scored on the PEDro scale, the final study assessment scores were defined. According to Maher et al. [29], a score between 8–11 is considered to be optimal, but if the study gains between 0–3 points, that study will be classified with “poor” quality, 4–5 points with “fair” quality, 6–8 points with “good” quality, and 9–10 points with “excellent” quality. Of all studies included in this systematic review, 2 studies showed fair quality, 17 of them showed good quality, and the other 2 studies showed excellent quality, which is shown in Table 1.

Table 1. PEDro scale results.

Study	Criterion											Σ
	1	2	3	4	5	6	7	8	9	10	11	
Jourkesh et al. (2011) [30]	Y	N	Y	N	Y	N	N	Y	Y	Y	Y	6
Kaminska et al. (2012) [31]	Y	N	Y	N	Y	N	N	Y	Y	Y	Y	6
Adriana et al. (2012) [32]	Y	N	Y	Y	Y	N	N	Y	Y	Y	Y	7
Ribeiro et al. (2013) [33]	Y	N	Y	N	Y	Y	N	Y	Y	Y	Y	7
Elamaram (2014) [34]	Y	Y	Y	N	Y	N	N	Y	Y	Y	Y	7
Çitozi et al. (2016) [35]	Y	N	Y	Y	N	N	N	Y	Y	Y	Y	6
Mitrović et al. (2016) [36]	Y	N	Y	Y	Y	N	N	Y	Y	Y	Y	7
Moskovljević (2016) [37]	Y	N	Y	Y	Y	N	N	Y	Y	Y	Y	7
Skurikhina et al. (2016) [38]	Y	N	N	Y	Y	N	N	Y	Y	Y	Y	6
Zou et al. (2016) [39]	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	9
Özcan et al. (2018) [40]	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	9
Suri et al. (2018) [41]	Y	N	N	Y	N	N	N	Y	Y	Y	Y	5
Pituk (2019) [42]	Y	N	N	Y	Y	N	N	Y	Y	Y	Y	6
Spiliopoulou et al. (2019) [43]	Y	N	N	Y	N	N	N	Y	Y	Y	Y	5
Wang et al. (2019) [44]	Y	N	Y	N	Y	N	N	Y	Y	Y	Y	6

Table 1. Cont.

Study	Criterion											Σ
	1	2	3	4	5	6	7	8	9	10	11	
Griban et al. (2020) [45]	Y	N	Y	N	Y	Y	N	Y	Y	Y	Y	7
Griban et al. (2020) [46]	Y	N	Y	N	Y	Y	N	Y	Y	Y	Y	7
Osipov et al. (2020) [47]	Y	N	N	N	Y	Y	N	Y	Y	Y	Y	6
Zhai et al. (2020) [48]	Y	N	N	N	Y	Y	N	Y	Y	Y	Y	6
Kang et al. (2021) [49]	Y	N	N	N	Y	Y	N	Y	Y	Y	Y	6
Shimamoto et al. (2021) [50]	Y	N	Y	N	Y	Y	N	N	Y	Y	Y	6

Legend: 1—eligibility criteria; 2—random allocation; 3—concealed allocation; 4—baseline comparability; 5—blind subject; 6—blind clinician; 7—blind assessor; 8—adequate follow-up; 9—intention-to-treat analysis; 10—between-group analysis; 11—point estimates and variability; Y—criterion is satisfied; N—criterion is not satisfied; Σ —total awarded points.

3.2. Selection and Characteristics of Studies

A search of electronic databases and scanning the reference lists yielded 11,839 studies. After removing duplicates, a total of 3938 studies were screened. Additional 3892 studies were excluded based on inclusion criteria and a total of 46 studies were screened and selected for eligibility. After increased sensitivity and in-deeper check, 25 studies with nonrelevant outcomes, editorials, and executive summaries were additionally excluded. Lastly, a total of 21 full-text studies were included in the systematic review (Figure 1).




Figure 1. Collecting adequate studies on the basis of pre-defined criteria (PRISMA flow chart).

Tables 2 and 3 show in more detail the studies that met the set conditions and entered the qualitative analysis.

Table 2. Review of studies.

First Author and Year of Publication	Sample of Participants			PF	Results
	Number	Age (Years)	Groups		
Jourkesh et al. (2011) [30]	N-450 M-250 F-200	M-22.5 $\pm$ 8.25 F-22.75 $\pm$ 6.20	FSPE	SaR, CMJ, 10 m SR, Flex, Pu	M > F (CMJ, 10 m SR, Flex, Pu) F > M (SaR)
Kaminska et al. (2012) [31]	N-82 M-28 F-54	19–23	FSPE, PH	Su, dyn, SB	** FSPE and PH
Adriana et al. (2012) [32]	X	X	(BuchUni) EG, CG, BB	30 m sprint, Su, Pu, CMJ	EG results are below average
Ribeiro et al. (2013) [33]	N-257 M-125 F-132	M-20.4 $\pm$ 2.3 F-19.9 $\pm$ 1.91	FSPE	MAT, SaR, 12 min run-walk test	High-level of PF Low level of CRP
Elamaran (2014) [34]	F-45	18–20	(IndUniP) ImT, ItT, CG	Sq, Pu, HJ, bdps, pnk	* ImT and ItT
Çitozi et al. (2016) [35]	N-24 M-14 F-10	19–21	FSPE	SB	SB increased by 16.08% (open eyes) and 20.62% (closed eyes)
Mitrović et al. (2016) [36]	N-137 M-67 F-70	19–21	KPA	dhg, mif (dl), LJ, HJ, Su, Plps, Pu	F (dhg = 6.99%, LJ = 4.59%, HJ = 7.3%) M (LJ = 2.83%, HJ = 4.27%)
Moskovljević (2016) [37]	N-58 M-29 F-29	20–21	FSPE	SB, Flex, Els, Rls	High level of PF students who have passed RG
Skurikhina et al. (2016) [38]	F-40	19–24	2 students groups	Pu, MAT, Rls	** PF
Zou et al. (2016) [39]	N-30 M-18 F-12	20.8 $\pm$ 2.07	(ChiUni) EG (Bd), CG	Ag, Bd, Els, LJ	** EG
Özcan et al. (2018) [40]	F-60	18–25	(FSPE) EG (AP), CG	SaR, dhg, dynb, dynl, DB, VC	** EG (AP)
Suri et al. (2018) [41]	F-96	19	FSPE	SaR, Bst	PEC ** Flex
Pituk (2019) [42]	N-392 M-167 F-225	M-18.4 $\pm$ 0.74 F-18.4 $\pm$ 1	FilUni	20 m SR, SaR, Z, Cu, CMJ	F > M (SaR) M > F (20 m SR, CMJ, Cu)
Spiliopoulou et al. (2019) [43]	F-20	21.8 $\pm$ 2.8	FSPE (ST, STIAT)	CMJ	STIAT showed better results in CMJ
Wang (2019) [44]	N-1414	18–24	TsgUni	LJ, VC, IFff-DA	Higher PA degree students have 2.39 $\times$ better scores at IFff-DA and 1.39 $\times$ at LJ
Griban et al. (2020) [45]	N-369 M-195 F-174	19–24	PolNatUni, 16 groups	100, 2000 and 3000 m sprint, LJ, Plps, Pu, MAT, 4 $\times$ 9 SR, SaR	Unsatisfactory level of PF

Table 2. Cont.

First Author and Year of Publication	Sample of Participants			PF	Results
	Number	Age (Years)	Groups		
Griban et al. (2020) [46]	N-394 M-199 F-195	19–24	PolNatUni, 16 groups	100, 2000 and 3000 m sprint, LJ, Plps, Pu, MAT, 4 × 9 SR, SaR	2nd year students show the best PF
Osipov et al. (2020) [47]	M-205 E1-127 E2-78	19–20	RusUniH, RusUniT	1mile-run test, Pu, sq, pnk	* of students who have the evidence of PA
Zhai et al. (2020) [48]	N-2324 M-1621 F-708	19.6 ± 0.6	3 ChiUni	VC, 50 m sprint, SaR, LJ, Su, Plps 1000 and 500 m run	PF ++ Agr no matter of LS
Kang et al. (2021) [49]	N-1183 M-832 F-351	23.2 ± 2.6	KorUni	VO2max on T	M > F VO2max
Shimamoto et al. (2021) [50]	N-95 M-52 F-33	18.9 ± 1.4	JapUni	VO2max on CE	VO2max = 42.5 ± 6.5 ML/kg/min

Legend: N—total number of participants; M—male, F—female; X—no data; EG—experimental group; CG—control group; PF—physical fitness tests; PA—physical activity; PEC—physical education classes; CMJ—countermovement jump; CRP—cardiorespiratory fitness; Els—explosive legs strength; Rls—repetitive legs strength; DB—dynamic balance; SB—static balance; mif—maximal isometric force; sq—squat; dhg—dynamometer hand grip; dynb—dynamometry back muscles; dynl—dynamometry legs; dl—deadlift; bdps—bench dips; KPA—Criminal Police Academy students; TsgUni—Tsinghua University students; PolNatUni—Poliand National University students; FilUni—Filipino University students; SPE—special physical education; ImT—isometric training; ItT—isotonic training; CRT—cardiorespiratory fitness; VO2max—maximal oxygen uptake; T—treadmill; CE—cycle ergometer; PH—physiotherapy students; FSPE—faculty of sport and physical education students; JapUni—Japanese University students; KorUni—Korean University students; ChiUni—Chinese University students; RusUniH—Russian University students of humanities specialties; RusUniT—Russian University students of technical specialties; BuchUni—Bucharest University; IndUniP—Indian University Polytechnic College; ST—strength training; STIAT—strength training and intensive aerobic training; Flex—flexibility; Pu—push-ups; Plps—pull-ups; Su—sit-ups; LJ—long jump; Z—zipper test; Cu—curl up test; HJ—high jump; IFff—isometric force of the finger flexor; DA—dominant arm; RG—rhythmic gymnastics classes; AP—aqua-pilates; sq—squat; pnk—plank; Agr—academic grades; MAT—modified abdomen test; SR—shuttle run; SaR—sit and reach; Bst—back scratch test; Bd—Bamby dance; Ag—agility; BB—basketball; VC—vital capacity; LS—lifestyle; * significant improvement; ** significant difference between groups; ++ positive correlation.

Table 3. Physical activity evaluation.

First Author and Year of Publication	Sample of Participants			PA Evaluation	Results
	Number	Age (Years)	Groups		
Kaminska et al. (2012) [31]	N-82 M-28 F-54	19–23	FSPE, PH	SAQ	Low level (M-10.7%, F-48.1%) Moderate level (M-35.7%, F-40.7%) High level (M-17.9%, F-5.6%) Vigorous level (M-35.7%, F-5.6%)
Pituk et al. (2019) [42]	N-392 M-167 F-225	M-18.4 ± 0.74 F-18.4 ± 1	FilUni	IPAQ	High level 37% Moderate level 48% Low level 15%
Wang et al. (2019) [44]	N-1414	18–24	TsgUni	CL-IPAQ	TA-41.50% LA-17.36% DA-7.5%

Table 3. *Cont.*

First Author and Year of Publication	Sample of Participants			PA Evaluation	Results
	Number	Age (Years)	Groups		
Osipov et al. (2020) [47]	M-205 E1-127 E2-78	19–20	RusUniH, RusUniT	FC, SS, IPAQ	* E1 in workplace PA * E2 in total PA
Zhai et al. (2020) [48]	N-2324 M-1621 F-703	19.6 ± 0.6	3 ChiUni	CS-IPAQ	** M in MVPA ** F in GPA
Kang et al. (2021) [49]	N-1183 M-832 F-351	23.2 ± 2.6	KorUni	Q	METs were gradual in order of PiA, FE and poor CRF
Shimamoto et al. (2021) [50]	N-95 M-52 F-33	18.9 ± 1.4	JapUni	Acc	EEPA and Ds were higher in part-time job students

Legend: N—total number of participants; M—male; F—female; CT—circuit training; A—aerobic; PH—physiotherapy students; FSPE—faculty of sport and physical education students; FilUni—Filipino University students; RusUniH—Russian University students of humanities specialties; RusUniT—Russian University students of technical specialties; ChiUni—Chinese University students; KorUni—Korean University students; JapUni—Japanese University students; TsgUni—Tsinghua University students; LA—learning activity; TA—traffic activity; DA—domestic work; IPAQ—International Physical Activity Questionnaire; SAQ—Self-Assessment Questionnaire; CL-IPAQ—Chinese long format of IPAQ; PA—physical activity; GPA—average grade point; MVPA—moderate-vigorous intensity physical activity; CS-IPAQ—Chinese short format of IPAQ; Q—questionnaire, BSIQ—Body Self Image Questionnaire; FC—fitness center documents; SS—sport school documents; PiA—physical inactivity; FE—fast eating; CRF—cardiorespiratory fitness; Acc—accelerometer; METs—metabolic equivalent; EEPA—energy exposure originating from physical activity per day; Ds—daily steps; * significant improvement; ** significant difference between groups.

There were a total of 7675 participants. The highest number was 2324 [48], the lowest was 20 [43], while for only one study there were no data about the total number of participants [32]. The sample of participants was only female in five studies [34,38,40,41,43], while only one study had male participants only [47]. There were eight studies where the sample of participants was from the Faculty of Sport and Physical Education students [30,31,33,35,37,40,41,43], only one study had Criminal Police Academy students [36], while the rest of the studies had participants from different universities and faculties [42–50].

For measuring physical fitness, it should be mentioned Criminal Police Academy students [36], who were using standard battery test, prescribed by plan and program “Special physical education”. Two studies had students from Poland National University [45,46], one had Tsingua University students from China [44], and all three mentioned were using their own standardized battery test. Aqua-Pilates program [40], Bamboo dance program [39], hiking activities [35], and fitness yoga [38] were also evaluated and there were also two studies that conducted VO2max evaluation on the treadmill [49] and cycle ergometer [50]. The rest of the studies were using standard and already known physical fitness tests. There were evaluations of physical fitness based on practicing basketball games [32] and rhythmic gymnastics, both conducted on faculty [37], and concurrent power training with high-intensity interval cycling [43].

Two types of relationships were conducted. The first was the relationship between physical fitness and academic performance among Chinese University students [48] and the second one [50] was a relationship between depression, daily physical activity, physical fitness, and daytime sleep time in Japanese University students. In addition, only one study [49] was determining predictors of metabolic syndrome (eating speed, physical activity and cardiorespiratory fitness) in Korean University students.

Evaluation of physical activity was various. International Physical Activity Questionnaire (IPAQ) was used in four studies [42,44,47,48] and among those mentioned, only Osipov et al. [47] were using fitness center and sport school documents as evaluation of the physical activity. The self-assessment questionnaire was conducted only

by Kaminska et al. [31], Kang et al. [49] had a regular questionnaire, while Shimamoto et al. [50] was the only study that monitored and evaluated physical activity using the accelerometer. Additionally, the mentioned questionnaires are valid and reliable [51–53], while the details of the accelerometer measurement have already been described in previous studies [54,55].

4. Discussion

The current study aimed to examine the scientific evidence regarding physical activity and physical fitness among university students. The universities are an ideal environment for the promotion of physical fitness and physical activity. The main findings of the current review are that university students show moderate levels of physical fitness and physical activity. However, when considering the results, it should be done with caution due to cultural differences, different faculties included in the review as well as the difference in educational systems. Moreover, some studies included students from the faculty of sports who have physical activities within the curriculum that must be taken into account. This was confirmed by Kaminska et al. [31] who found that students of the faculty of sports have better physical fitness compared to physiotherapy students, which was explained by a higher degree of physical activity in classes.

Additionally, when it comes to analyzing the impact of physical activities, such as aqua-Pilates [40], it was concluded that the group that implemented the program made statistically significant progress in terms of muscle strength, flexibility and balance. Meredith-Jones et al. [56] have also found similar results, that water exercises can lead to beneficial effects on cardiorespiratory fitness, strength and body fat. The weight loss in water removes the body load and helps in increasing the range of motion according to normal conditions, as well as significant energy expenditure [57]. In addition, such results are a consequence of the application of exercises that develop the mentioned abilities, because as already mentioned, the application of physical activities can increase the already acquired level of fitness and physical abilities. Fitness is an activity that consists of exercises that require the engagement of the whole body and develop both coordination and strength, as well as flexibility and balance [38].

Jourkesh et al. [30] have found that females performed better in flexibility tests than males, which is in full accordance with Ortega et al. [58]. Hiking activities conducted by Citozi et al. [35] found gender differences in balance tests, which is lined with other findings [59,60]. In accordance to previously mentioned studies, Zou et al. [39] have examined the influence of Bamboo dance and at the end of the program, the experimental group showed significant progress in balance (3.6%), agility (0.18%), strength (0.33%) and explosive power (0.42%). As the authors have considered low physical activity in university students, Bamboo dance interventions are needed. According to all mentioned facts, the main focus should be placed on reducing barriers that students experience that may impact their physical activity [61].

Wang [44] results showed that students with a higher degree of physical activity have 2.39 times better results in the strength test and 1.39 times in the long jump test. Osipov et al. [47] also used a standardized questionnaire to assess physical activity to divide participants into groups, and proved that participants with a higher degree of physical activity had better physical fitness compared to those with a lower degree. The results were even compared with the obtained results of some other countries, where Russian students show significantly better results compared to students from African countries [62,63], Turkey [64], Iran [65] and Ukraine [66], as well as several more European countries [67,68]. Two similar studies conducted by Griban et al. [45,46] gave two new answers. The first indicates the problem of increasing the level of fitness in higher institutions and the second is consisted of analyzing the dynamics of fitness on the same sample. Likewise, Adriana et al. [32] found that younger students have poorer results in physical fitness compared to older ones, in exact same variables. Suri et al. [41] are suggesting that there is a strong need for more active physical education programs that are appropriate for

developing fitness and improving the health status of college-going students. Although the current state of physical fitness of these students was at a very unsatisfactory level, the authors emphasize that the system primarily does not provide the required level of physical fitness and work ability, which is why there is a great need to identify an adequate and efficient program that will improve the traditional physical education system. Kang et al. [49] found physical inactivity and poor cardiorespiratory function to metabolic equivalent of task risk. Students who are busy with their studies, part-time jobs and extracurricular sport clubs—physical activities that are normally considered to be beneficial [69,70] may cause excessive negative side effects [50]. In accordance with the current findings, similar results have been found in Western [71–74] and Asian countries [75–77].

Unlike all other studies in which the authors wanted to show that applying physical activity has a positive impact on the physical fitness of students, Mitrović et al. [36] decided to examine the impact of the absence of special physical education on physical fitness in an 8-month period. They came to the conclusion that the absence of this type of physical activity in students causes a significant decline in physical fitness, which may be an indication that already acquired physical fitness may decline if physical activity is not practiced. If there is more physical activity, the more physically fit individuals at the adequate level are there [78]. According to Rossomanno et al. [79], if we want to secure that these students could perform their duties, it is necessary to apply the monitored program for the development of physical fitness and physical activity during the whole year.

The main limitation of this study lies in cultural differences, where we included a wide range of universities, as well as differences in educational systems. As studies about physical activity interventions have rarely been enforced in some parts of included studies, further research should promote regular exercise in the university students. Despite the mentioned limitations, this study should be an important contribution to physical activity, as well as physical fitness research, and be beneficial to elucidating the primary negative factors.

5. Conclusions

Based on the obtained results, university students show a satisfactory level of physical activity and physical fitness. However, the results vary due to the different factors involved, which are mostly related to cultural differences and educational systems in different countries. The results for the students of the faculty of sport showed that physical activity has a positive effect on the development and maintenance of physical fitness and activity. As far as students from different faculties, this study suggests that further investigations should be conducted to promote daily exercise, as it may be beneficial for physical fitness and activity related to students health.

Author Contributions: Conceptualization, V.K. and G.S.; methodology, V.K. and G.S.; software, M.O.; validation, K.P. and D.T.-P.; formal analysis, M.O. and R.J.; investigation, D.Đ.; resources, M.S. and D.Đ.; data curation, M.Č.; writing—original draft preparation, D.Đ.; writing—review and editing, D.T.-P. and K.P.; visualization, M.Č.; supervision, D.T.-P., R.J. and K.P.; project administration, M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Strong, W.B.; Malina, R.M.; Blimkie, C.J.R.; Daniels, S.R.; Dishman, R.K.; Gutin, B.; Hergenroeder, A.C.; Must, A.; Nixon, P.A.; Pivarnik, J.M. Evidence based physical activity for school-age youth. *J. Pediatr.* **2005**, *146*, 732–737. [CrossRef]
- Warburton, D.E.R.; Nicol, C.W.; Bredin, S.S.D. Health benefits of physical activity: The evidence. *CMAJ* **2006**, *174*, 801–809. [CrossRef]
- Janssen, I.; LeBlanc, A.G. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int. J. Behav. Nutr. Phys. Act.* **2010**, *7*, 1–16. [CrossRef] [PubMed]
- Petrović-Oggiano, G.P.; Damjanov, V.; Gurinović, M.A.; Glibetić, M.D. Physical activity in prevention and reduction of cardiovascular risk. *Med. Pregl.* **2010**, *63*, 200–207. [CrossRef] [PubMed]
- Ortega, F.B.; Ruiz, J.R.; Castillo, M.J. Physical activity, physical fitness, and overweight in children and adolescents: Evidence from epidemiologic studies. *Endocrinol. Nutr. Engl. Ed.* **2013**, *60*, 458–469. [CrossRef] [PubMed]
- Diethelm, K.; Huybrechts, I.; Moreno, L.; De Henauw, S.; Manios, Y.; Beghin, L.; Gonzalez-Gross, M.; Le Donne, C.; Cuenca-Garcia, M.; Castillo, M.J. Nutrient intake of European adolescents: Results of the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Publ. Health Nutr.* **2014**, *17*, 486–497. [CrossRef] [PubMed]
- Mitić, D. Značaj fizičke aktivnosti u prevenciji i terapiji gojaznosti u detinjstvu i adolescenciji. *Med. Glas. Spec. Boln. Boles Štitaste žlezde Boles. Metab.* **2011**, *16*, 107–112.
- Mandarić, S. Effects of programmed exercising to music of female pupils. *Facta Univ. Phys. Educ. Sport* **2001**, *1*, 37–49.
- Bouchard, C.; Després, J.-P. Physical activity and health: Atherosclerotic, metabolic, and hypertensive diseases. *Res. Q. Exerc. Sport* **1995**, *66*, 268–275. [CrossRef] [PubMed]
- U.S. Department of Health and Human Services. Physical Activity and Health: A Report of the Surgeon General. Natl Cent Chronic Dis Prev Heal Promot. Available online: <https://www.cdc.gov/nccdphp/sgr/index.htm> (accessed on 19 November 2021).
- Schipperijn, J.; Cerin, E.; Adams, M.A.; Reis, R.; Smith, G.; Cain, K.; Christiansen, L.B.; Van Dyck, D.; Gidlow, C.; Frank, L.D. Access to parks and physical activity: An eight country comparison. *Urban For. Urban Green.* **2017**, *27*, 253–263. [CrossRef] [PubMed]
- Sallis, J.F.; Conway, T.L.; Cain, K.L.; Carlson, J.A.; Frank, L.D.; Kerr, J.; Glanz, K.; Chapman, J.E.; Saelens, B.E. Neighborhood built environment and socioeconomic status in relation to physical activity, sedentary behavior, and weight status of adolescents. *Prev. Med.* **2018**, *110*, 47–54. [CrossRef] [PubMed]
- Franck, C.; Grandi, S.M.; Eisenberg, M.J. Taxing junk food to counter obesity. *Am. J. Public Health* **2013**, *103*, 1949–1953. [CrossRef] [PubMed]
- Kohn, M.; Golden, N.H. Eating disorders in children and adolescents. *Paediatr. Drugs* **2001**, *3*, 91–99. [CrossRef] [PubMed]
- Caia, J.; Weiss, L.W.; Chiu, L.Z.F.; Schilling, B.K.; Paquette, M.R.; Relyea, G.E. Do lower-body dimensions and body composition explain vertical jump ability? *J. Strength Cond. Res.* **2016**, *30*, 3073–3083. [CrossRef] [PubMed]
- Kubieva, S.S.; Botagariev, T.A.; Konisbaeva, S.I.; Mambetov, N.; Aralbayev, A.S.; Zhetimekov, Y.T.; Dairabayev, S.E.; Kaliyakbarova, S.K.; Sadykova, Z.L. Health promotion program for the students with regard to the level of their physical activity, physical fitness and health. *J. Phys. Educ. Sport* **2019**, *19*, 703–709.
- DeBar, L.L.; Schneider, M.; Drews, K.L.; Ford, E.G.; Stadler, D.D.; Moe, E.L.; White, M.; Hernandez, A.E.; Solomon, S.; Jessup, A. Student public commitment in a school-based diabetes prevention project: Impact on physical health and health behavior. *BMC Publ. Health* **2011**, *11*, 1–11. [CrossRef] [PubMed]
- Andersen, L.B.; Harro, M.; Sardinha, L.B.; Froberg, K.; Ekelund, U.; Brage, S.; Anderssen, S.A. Physical activity and clustered cardiovascular risk in children: A cross-sectional study (The European Youth Heart Study). *Lancet* **2006**, *368*, 299–304. [CrossRef]
- Kwan, M.Y.; Cairney, J.; Faulkner, G.E.; Pullenayegum, E.E. Physical activity and other health-risk behaviors during the transition into early adulthood: A longitudinal cohort study. *Am. J. Prev. Med.* **2012**, *42*, 14–20. [CrossRef] [PubMed]
- Bray, S.R.; Born, H.A. Transition to university and vigorous physical activity: Implications for health and psychological well-being. *J. Am. Coll. Health* **2004**, *52*, 181–188. [CrossRef]
- Crombie, A.P.; Ilich, J.Z.; Dutton, G.R.; Panton, L.B.; Abood, D.A. The freshman weight gain phenomenon revisited. *Nutr. Rev.* **2009**, *67*, 83–94. [CrossRef] [PubMed]
- Vella-Zarb, R.A.; Elgar, F.J. The ‘freshman 5’: A meta-analysis of weight gain in the freshman year of college. *J. Am. Coll. Health* **2009**, *58*, 161–166. [CrossRef]
- Suminski, R.R.; Petosa, R.; Utter, A.C.; Zhang, J.J. Physical activity among ethnically diverse college students. *J. Am. Coll. Health* **2002**, *51*, 75–80. [CrossRef] [PubMed]
- Vuillemin, A.; Boini, S.; Bertrais, S.; Tessier, S.; Oppert, J.-M.; Hercberg, S.; Guillemin, F.; Briançon, S. Leisure time physical activity and health-related quality of life. *Prev. Med.* **2005**, *41*, 562–569. [CrossRef] [PubMed]
- DeVahl, J.; King, R.; Williamson, J.W. Academic incentives for students can increase participation in and effectiveness of a physical activity program. *J. Am. Coll. Health* **2005**, *53*, 295–298. [CrossRef] [PubMed]
- Arzu, D.; Tuzun, E.H.; Eker, L. Perceived barriers to physical activity in university students. *J. Sports Sci. Med.* **2006**, *5*, 615. [PubMed]
- Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G.; Group, P. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* **2009**, *6*, e1000097. [CrossRef]

28. de Morton, N.A. The PEDro scale is a valid measure of the methodological quality of clinical trials: A demographic study. *Aust. J. Physiother.* **2009**, *55*, 129–133. [\[CrossRef\]](#)
29. Maher, C.G.; Sherrington, C.; Herbert, R.D.; Moseley, A.M.; Elkins, M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys. Ther.* **2003**, *83*, 713–721. [\[CrossRef\]](#)
30. Jourkesh, M.; Sadri, I.; Ojagi, A.; Sharanavard, A. Determination of fitness level in male and female college aged students. *Arch. Appl. Sci. Res.* **2011**, *3*, 326–333.
31. Irena, K.; Anna, M.; Anda, B. Physical activity and its relation to health-related physical fitness in students. *Ovidius Univ. Ann. Ser. Phys. Educ. Sport Mov. Health* **2012**, *12*, 256–264.
32. Adriana, S.; Mihaela, G. Physical ability of the students practicing basketball games-Comparative study. *Ovidius Univ. Ann. Ser. Phys. Educ. Sport Mov. Health* **2012**, *12*, 227–233.
33. Ribeiro, A.; Silva, D.; Carvalho, F.; Schiavoni, D.; Jesus, B.; Cyrino, E. Health-related physical fitness in men and women aged 17–26 years. *Rev. Bras. Ativid. Física Saúde* **2013**, *18*, 197. [\[CrossRef\]](#)
34. Elamaram, M. Effect of different methods of strength training on selected motor fitness components of women college students. *Int. J. Inf. Res. Rev.* **2014**, *1*, 173–175.
35. Çitozi, R.; Kasa, A.; Peja, E. The role of hiking activities in improving balance skills to the students of Sports University of Tirana. *Arena J. Phys. Act.* **2016**, *5*, 69–78.
36. Mitrović, B.; Janković, R.; Dopsaj, M.; Vučković, G.; Milojević, S.; Pantelić, S.; Nurkić, M. How an eight-month period without specialized physical education classes affects the morphological characteristics and motor abilities of students of the Academy of Criminalistic and Police Studies. *Facta Univ. Phys. Educ. Sport* **2016**, *14*, 167–178.
37. Moskovljević, L. The relations between certain motor abilities and success in rhythmic gymnastics in the students of different genders. *Fizička Kult.* **2016**, *70*, 155–163. [\[CrossRef\]](#)
38. Skurikhina, N.V.; Kudryavtsev, M.D.; Kuzmin, V.A.; Iermakov, S. Fitness yoga as modern technology of special health groups' girl students' psycho-physical condition and psycho-social health strengthening. *Phys. Educ. Stud.* **2016**, *2*, 24–31. [\[CrossRef\]](#)
39. Zou, L.; Zeng, N.; He, J. The effect of bamboo dance on motor fitness in Chinese college students. *Int. J. Phys. Educ. Sport Health* **2016**, *3*, 193–199.
40. Özcan, R.; İrez, G.B.; Saygin, Ö.; Ceylan, H.İ. Aqua-pilates exercises improves some physical fitness parameters on healthy young women. *J. Phys. Educ. Sport Sci. Egit. Sport Bilim. Derg.* **2018**, *12*, 160–175.
41. Suri, M.; Sharma, R.; Saini, N. Effect of physical education and physical activity on anthropometric measurements and flexibility among college going girls. *Eur. J. Phys. Educ. Sport Sci.* **2018**, *4*, 62–72.
42. Pituk, C.S.; Cagas, J.Y. Physical Activity and Physical Fitness among Filipino University Students. *J. Phys. Educ.* **2019**, *30*, 1–10.
43. Spiliopoulou, P.; Zaras, N.; Methenitis, S.; Papadimas, G.; Papadopoulos, C.; Bogdanis, G.C.; Terzis, G. Effect of concurrent power training and high-intensity interval cycling on muscle morphology and performance. *J. Strength Cond. Res.* **2019**, *35*, 2464–2471. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Wang, J. The association between physical fitness and physical activity among Chinese college students. *J. Am. Coll. Health* **2019**, *67*, 602–609. [\[CrossRef\]](#)
45. Griban, G.; Yahupov, V.; Svystun, V.; Dovgan, N.; Yeromenko, E.; Udych, Z.; Zhuravlov, I.; Kushniriuk, S.; Semeniv, B.; Konoval'ska, L. Dynamics of the students' physical fitness while studying at higher educational institutions. *Int. J. Appl. Exerc. Physiol.* **2020**, *9*, 147–156.
46. Griban, G.; Dovgan, N.; Tamozhanska, G.; Semeniv, B.; Ostapenko, A.; Honcharuk, N.; Khurtenko, O.; Kozibroda, L.; Husarevych, O.; Denysovets, A. State of physical fitness of the students of Ukrainian higher educational institutions. *Int. J. Appl. Exerc. Physiol.* **2020**, *9*, 16–26.
47. Osipov, A.Y.; Potop, V.; Nagovitsyn, R.S.; Zemba, E.A.; Knyazev, A.P.; Orlova, I.I.; Ratmanskaya, T.I.; Iermakov, S.S. Indicators of physical activity and fitness of male students at Russian universities. *Phys. Educ. Stud.* **2020**, *24*, 40–46. [\[CrossRef\]](#)
48. Zhai, X.; Ye, M.; Gu, Q.; Huang, T.; Wang, K.; Chen, Z.; Fan, X. The relationship between physical fitness and academic performance among Chinese college students. *J. Am. Coll. Health* **2020**, *5*, 1–9. [\[CrossRef\]](#)
49. Kang, M.; Joo, M.; Hong, H.; Kang, H. Eating Speed, Physical Activity, and Cardiorespiratory Fitness Are Independent Predictors of Metabolic Syndrome in Korean University Students. *Nutrients* **2021**, *13*, 2420. [\[CrossRef\]](#)
50. Shimamoto, H.; Suwa, M.; Mizuno, K. Relationships between depression, daily physical activity, physical fitness, and daytime sleepiness among Japanese university students. *Int. J. Environ. Res. Public Health* **2021**, *18*, 8036. [\[CrossRef\]](#)
51. Rowe, D.A.; Benson, J.; Baumgartner, T.A. Development of the body self-image questionnaire. *Meas. Phys. Educ. Exerc. Sci.* **1999**, *3*, 223–247. [\[CrossRef\]](#)
52. Dinger, M.K.; Behrens, T.K.; Han, J.L. Validity and reliability of the International Physical Activity Questionnaire in college students. *Am. J. Health Educ.* **2006**, *37*, 337–343. [\[CrossRef\]](#)
53. Sasaki, S.; Katagiri, A.; Tsuji, T.; Shimoda, T.; Amano, K. Self-reported rate of eating correlates with body mass index in 18-y-old Japanese women. *Int. J. Obes.* **2003**, *27*, 1405–1410. [\[CrossRef\]](#)
54. Ayabe, M.; Aoki, J.; Kumahara, H.; Tanaka, H. Assessment of minute-by-minute stepping rate of physical activity under free-living conditions in female adults. *Gait Posture* **2011**, *34*, 292–294. [\[CrossRef\]](#) [\[PubMed\]](#)
55. Aoyagi, Y.; Shephard, R.J. Habitual physical activity and health in the elderly: The Nakanojo Study. *Geriatr. Gerontol. Int.* **2010**, *10*, S236–S243. [\[CrossRef\]](#) [\[PubMed\]](#)

56. Meredith-Jones, K.; Waters, D.; Legge, M.; Jones, L. Upright water-based exercise to improve cardiovascular and metabolic health: A qualitative review. *Complementary Ther. Med.* **2011**, *19*, 93–103. [[CrossRef](#)]
57. Kantyka, J.; Herman, D.; Rocznio, R.; Kuba, L. Effects of aqua aerobics on body composition, body mass, lipid profile, and blood count in middle-aged sedentary women. *Hum. Mov.* **2015**, *16*, 9–14. [[CrossRef](#)]
58. Ortega, F.B.; Artero, E.G.; Ruiz, J.R.; España-Romero, V.; Jiménez-Pavón, D.; Vicente-Rodríguez, G.; Moreno, L.A.; Manios, Y.; Beghin, L.; Ottevaere, C. Physical fitness levels among European adolescents: The HELENA study. *Br. J. Sport Med.* **2011**, *45*, 20–29. [[CrossRef](#)]
59. Wieczorek, A.; Adrian, J. The level of motor abilities in consideration of health conditions of selected youth groups. *Stud. Phys. Cult. Tour.* **2006**, *13*, 85–92.
60. Toole, T.; Kretschmar, J.C. Gender differences in motor performance in early childhood and later adulthood. *Women Sport Phys. Act. J.* **1993**, *2*, 41–71. [[CrossRef](#)]
61. Roberts, S.; Reeves, M.; Ryrrie, A. The influence of physical activity, sport and exercise motives among UK-based university students. *J. Furth. High. Educ.* **2015**, *39*, 598–607. [[CrossRef](#)]
62. Essaw, E.; Moses, M.O.; Afrifa, D.; Acheampong, I.K.; Mensah, W.; Owusu, L. Physical activity patterns and dietary habits of undergraduate students. *Balt. J. Health Phys. Act.* **2019**, *11*, 115–123. [[CrossRef](#)]
63. Tinazci, C.; EAlrefai, S.; Musa, O. Patterns of physical activity of Libyan undergraduate students at the university of Tripoli using international physical activity questionnaire (IPAQ). *Sport Mont.* **2019**, *17*, 103–106.
64. Bednarek, J.; Pomykała, S.; Bigosińska, M.; Szygula, Z. Physical activity of Polish and Turkish university students as assessed by IPAQ. *Cent. Eur. J. Sport Sci. Med.* **2016**, *16*, 13–22. [[CrossRef](#)]
65. Heravi, M.D.; Moeini, B.; Hazavehei, M.M.; Moghimbeigi, A.; Homayonfar, S.; Moslem, A.; Mirmoosavi, S.J. Relationship between blood pressure and physical activity in adults 20 to 65 years old. *Amaz. Investig.* **2018**, *7*, 285–294.
66. Bergier, J.; Tsos, A.; Popovych, D.; Bergier, B.; Niżnikowska, E.; Ács, P.; Junger, J.; Salonna, F. Level of and factors determining physical activity in students in Ukraine and the Visegrad countries. *Int. J. Environ. Res. Public Health* **2018**, *15*, 1738. [[CrossRef](#)]
67. Bergier, B.; Bergier, J.; Niznikowska, E.; Junger, J.; Salonna, F.; Acs, F.; Fromel, K. Differences in physical activity and nutrition and silhouette-related behaviours in male and female students in selected European countries. *Ann. Agric. Environ. Med.* **2018**, *25*, 176–181. [[CrossRef](#)]
68. Jagiello, W. Differentiation of the body composition in taekwondo-ITF competitors of the men's Polish national team and direct based athletes. *Arch. Budo* **2015**, *11*, 329–338.
69. Paluska, S.A.; Schwenk, T.L. Physical activity and mental health. *Sport Med.* **2000**, *29*, 167–180. [[CrossRef](#)]
70. Peluso, M.A.M.; de Andrade, L.H.S.G. Physical activity and mental health: The association between exercise and mood. *Clinics* **2005**, *60*, 61–70. [[CrossRef](#)]
71. Miller, J.M.; Street, B.D. Metabolic syndrome and physical activity levels in college students. *Metab. Syndr. Relat. Disord.* **2019**, *17*, 431–435. [[CrossRef](#)]
72. López-Martínez, S.; Sánchez-López, M.; Solera-Martínez, M.; Arias-Palencia, N.; Fuentes-Chacón, R.M.; Martínez-Vizcaíno, V. Physical activity, fitness, and metabolic syndrome in young adults. *Int. J. Sport Nutr. Exerc. Metab.* **2013**, *23*, 312–321. [[CrossRef](#)] [[PubMed](#)]
73. Díez-Fernández, A.; Martínez-Vizcaíno, V.; Torres-Costoso, A.; Cañete García-Prieto, J.; Franquelo-Morales, P.; Sánchez-López, M. Strength and cardiometabolic risk in young adults: The mediator role of aerobic fitness and waist circumference. *Scand. J. Med. Sci. Sport* **2018**, *28*, 1801–1807. [[CrossRef](#)] [[PubMed](#)]
74. Salonen, M.K.; Wasenius, N.; Kajantie, E.; Lano, A.; Lahti, J.; Heinonen, K.; Räikkönen, K.; Eriksson, J.G. Physical activity, body composition and metabolic syndrome in young adults. *PLoS ONE* **2015**, *10*, e0126737.
75. Lee, J.; Kim, S.-U.; Kang, H.-S. Low cardio/respiratory fitness as an independent predictor of metabolic syndrome in Korean young men. *Eur. J. Appl. Physiol.* **2010**, *108*, 633–639. [[CrossRef](#)]
76. Kim, D.; Kim, J.Y.; Lee, M.K.; Lee, H.-D.; Lee, J.-W.; Jeon, J.Y. The relationship between fitness, BMI and risk factors of metabolic syndrome among university students in Korea. *Korean J. Obes.* **2012**, *21*, 99–107. [[CrossRef](#)]
77. Ahn, S.-H.; Um, Y.-J.; Kim, Y.-J.; Kim, H.-J.; Oh, S.-W.; Lee, C.M.; Kwon, H.; Joh, H.-K. Association between physical activity levels and physical symptoms or illness among university students in Korea. *Korean J. Fam. Med.* **2016**, *37*, 279. [[CrossRef](#)] [[PubMed](#)]
78. Collingwood, T.R. Physical fitness leadership in law enforcement. *Pol. Chief* **1988**, *55*, 30–37.
79. Rossomanno, C.I.; Herrick, J.E.; Kirk, S.M.; Kirk, E.P. A 6-month supervised employer-based minimal exercise program for police officers improves fitness. *J. Strength Cond. Res.* **2012**, *26*, 2338–2344. [[CrossRef](#)]